



MP Biomedicals, LLC.

29525 Fountain Parkway
Solon, Ohio 44139

Telephone: 440/337-1200

Toll Free: 800/854-0530

Fax: 440/337-1180

mailto:biotech@mpbio.com

web: www.mpbio.com

cappel

**RABBIT IGG FRACTION TO
HAMSTER IGG (WHOLE MOLECULE) - 5ML**

Catalog #: 55398

Lot #: 00457

Total Protein: 12.0 mg/ml
Antibody Protein: 3 mg/ml

DESCRIPTION:

Product is the lyophilized powder of rabbit IgG fraction to hamster IgG (whole molecule) and buffer salts.

The IgG fraction is prepared from the specific rabbit antiserum by delipidation, ammonium sulfate fractionation and anion exchange chromatography. The resulting IgG fraction is dialyzed into 0.02M sodium phosphate, 0.14M sodium chloride, pH 7.3, adjusted to standard titer, filtered through a 0.22 μ m filter, vialled and lyophilized. No preservative is added.

USES:

This product is suitable for use as a precipitating antibody. It gives sharp precipitin lines in immunoelectrophoresis and immunodiffusion techniques.

This product is suitable for conjugation with enzymes, radiolabels, fluorochromes or other markers, and for attachment to solid supports for use as an immunoabsorbant.

This product is suitable for use as a primary reagent in enzyme immunoassays, fluorescein immunoassays and blot, cell or tissue immunostaining. See **Related Product Ordering Information**.

SUGGESTED WORKING DILUTIONS

1:50 - 1:200 for Tissue Staining
1:100 - 1:500 for Immunoblotting
1:400 - 2,000 for Elisa

It is recommended that each lab obtain their own optimal working dilution for each assay

SPECIFICATIONS:

The total protein is measured using the Biuret procedure, with bovine albumin as a standard. Antibody titer is standardized to an in-house standard by equivalence-point precipitation. Each IgG fraction is tested for purity and specificity at a minimum of 40 mg/ml using immunoelectrophoresis. The product is mostly rabbit IgG; no trace of albumin is detected. It shows reactivity to hamster IgG; cross-reactivity to other species may exist. Because the product is directed against whole IgG, some antibodies are expected to bind light chain sites common to all immunoglobulins. Antibody activity to non-immunoglobulin serum proteins is not present.

RECONSTITUTION:

Reconstitute product with 5.0 ml of deionized or distilled water. Gentle swirling may be used to speed rehydration. Avoid vigorous shaking of the reconstituted material.

STORAGE:

Store lyophilized product at 2-8°C. Store reconstituted material no longer than 1-2 days at 2-8°C, unless preservative is added. For long term storage, aliquot and store reconstituted product frozen at -20°C or lower.

WARNING AND CAUTION:

This product is intended **FOR RESEARCH USE ONLY**, not for use in diagnostic or therapeutic procedures involving humans or animals.

RELATED PRODUCT ORDERING INFORMATION:

Catalog # Description Size

55395 (01080082) RABBIT ANTI-HAMSTER IGG (WHOLE MOLECULE) 2ML
55401 (12080082) FITC RABBIT ANTI-HAMSTER IGG (WHOLE MOLECULE) 2ML
55933 (60080080) PURIFIED HAMSTER IGG 20MG


Approved by Joseph Dietz, Ph.D.
Quality Control Director

**TO PLACE AN ORDER IN THE U.S., PLEASE CALL 800-854-0530
OR TO OBTAIN TECHNICAL ASSISTANCE IN THE U.S., PLEASE CALL 800-279-5490
OUTSIDE THE U.S., CONTACT YOUR LOCAL DISTRIBUTOR.**