


TECHNICAL INFORMATION

Catalog Number: 193918

Etoposide

Structure:



Molecular Formula: C₂₉H₃₂O₁₃

Formula Weight: 588.57

CAS # 33419-42-0

Synonyms: 9-[(4,6-O-Ethylidene-β-D-glucopyranosyl)oxy]-5,8,8a,9-tetrahydro-5-(4-hydroxy-3,5-dimethoxyphenyl)furo[3',4':6,7]naphtho[2,3-d]-1,3-dioxol-6(5aH)-one; 4'-demethylepipodophyllotoxin
9-[4,6-O-ethylidene-β-D-glucopyranoside]

Solubility: Soluble in DMSO (30 mg/ml), acetonitrile (2 mg/ml) or in a 9:1 mixture of chloroform:methanol (5mg/ml).

References:

- Merck Index, 12th Ed., # 3931
- C. Keller-Jusln et al., U.S. pat. 3,524,844 (1970 to Sandoz)
- C. Keller-Jusln et al., *J. Med. Chem.*, v. 14, 936 (1971).
- H. Sthelin, *Eur. J. Cancer*, v. 9, 215 (1973).
- S. B. Horwitz, J. D. Loike, *Lloydia*, v. 40, 82 (1977).
- F. R. Pelsor et al., *J. Pharm. Sci.*, v. 67, 1106 (1978).
- R. L. Chard et al., *Cancer Treat. Rep.*, v. 63, 1755 (1979)
- R. E. Slayton et al., *ibid.* 2089.
- S. M. Sieber et al., *Teratology*, v. 18, 31 (1978).
- B. F. Issell, S. T. Crooke, *Cancer Treat. Rev.*, v. 6, 107-124 (1979).
- P. J. O'Dwyer et al., *N. Engl. J. Med.*, v. 312, 692-700 (1985).
- P. I. Clark, M. L. Slevin, *Clin. Pharmacokinet.*, v. 12, 223-252 (1987).
- J. J. M. Holthuis et al. in *Analytical Profiles of Drug Substances* vol. 18, K. Florey, Ed. (Academic Press, New York, 1989) pp 121-151.